

Childhood Immunizations Report

Indicator: Percentage of children who have received appropriate immunizations by their 2nd birthday during the measurement period.

Numerator: Children who have evidence showing they received recommended vaccines, had documented history of the illness, or had an allergic reaction to the vaccine by their second birthday.

- 4 DTaP Vaccines, each at least one day apart, between 42 days and 2 years after birth OR Anaphylactic Reaction to DTaP Vaccine, OR Encephalopathy within 2 years of birth
- 3 inactivated polio vaccines (IPV), each at least one day apart, between 42 days and 2 years after birth OR Anaphylactic Reaction to IPV, Streptomycin, Polymyxin, or Neomycin within 2 years of birth
- 1 measles, mumps and rubella (MMR) vaccine OR Disorders of the Immune System, HIV, Malignant Neoplasm of Lymphatic Tissue, OR Anaphylactic Reaction to Neomycin within 2 years of birth
- 3 Hemophilus Influenza B (HiB) Vaccines, each at least one day apart, between 42 days and 2 years after birth OR Anaphylactic Reaction to HiB within 2 years of birth
- 3 Hepatitis B Vaccines, each at least one day apart, within 2 years of birth OR Anaphylactic Reaction to Hepatitis B Vaccine or to Common Baker's Yeast within 2 years of birth OR A diagnosis (active or inactive) of Hepatitis B
- 1 Varicella Zoster Vaccine (VZV) OR Disorder of the Immune System, HIV, Malignant Neoplasm of Lymphatic Tissue, OR Anaphylactic Reaction to Neomycin
- 4 Pneumococcal Conjugate Vaccines, each at least one day apart, between 42 days and 2 years after birth OR Anaphylactic reaction to Pneumococcal Conjugate Vaccine within 2 years of birth
- 1 Hepatitis A Vaccine within 2 years of birth OR Anaphylactic Reaction to Hepatitis A Vaccine within 2 years of birth OR Hepatitis A
- 2 2-dose Rotavirus Vaccines, each at least one day apart, between 42 days and 2 years after birth OR 3 3-dose Rotavirus Vaccines, each at least one day apart, between 42 days and 2 years after birth OR Anaphylactic Reaction to Rotavirus Vaccine
- 2 Influenza Vaccines, each at least one day apart, between 180 days and 2 years after birth OR Anaphylactic Reaction to Influenza Vaccine, Disorders of the Immune System, HIV, Malignant Neoplasm of Lymphatic Tissue, OR Anaphylactic Reactions to Neomycin with 2 years of birth

Denominator: Children who turn 2 years of age at the end of the reporting period and who have a visit during the reporting period.

Exclusions: Hospice Care

Community Health Northwest Florida's Goal: 42.80% (National Standard)